

# **Post-Operative Care: Cubital Tunnel Release**

### Wound Care

- Suture removal at 2-3 weeks.

 Once sutures are removed, you are now able to get your elbow wet in the shower, hand washing, and use hand sanitizer.

- No soaking in contaminated water such as dirty dish water, pools, hot tubs, and lakes until surgical site is fully healed (2-3 weeks after the sutures are removed).

- Cover with bandage if you are going to be in a dirty environment. Otherwise, open to air.

## Limitations

 The first two weeks there is a weight limit for your arm of approximately 1 pound. Leave post-operative dressing on until post-op visit with clinic providers or therapy.

- Week 2-4 post-operatively the weight limit is no more than 5 pounds.

- Week 4-6 post-operatively you may start gradually to wean back into normal activity with pain being your guide.

- At week 6, there are no limitations, but remember, you will be sore with some activities.

# **Surgical Expectations**

- You will be sore over the surgical site for up to 3 months, especially with direct pressure.

- Soreness and swelling will fluctuate the first three months as you wean back into normal activity, this is normal.
- Numbness or decreased sensation around the surgical site is normal for a few months after surgery.

-Strength in the arm will also take time to return, be patient.

 Numbness and tingling can continue to linger but improve over 12 to 18 months, sometimes 2 years depending on the severity of your cubital tunnel syndrome.

#### Therapy

- You will attend therapy at least 1 to 2 times. You may need more therapy depending on how your recovery is going.

 Do the therapy exercises at home at least once a day for 6 weeks focusing on range of motion and scar massage.

- Therapy is key to a successful recovery.

#### Brace

- Use as much and for as long as needed if it does not affect range of motion.

- You can use an elbow pad off and on after surgery.

- Anytime you have a question.

When to Call 605.217.2667

- Decreased range of motion.

- Increased pain, redness, drainage, or numbness/tingling.